

27207 Lahser Road #200-B  
Southfield, MI 48034  
Ph. (248)663-1900

**Kyle Anderson, MD**  
**William Beaumont Hospital**  
6900 Orchard Lake Rd #103  
W. Bloomfield, MI 48322  
Ph. (248)855-7400

3535 W. 13 Mile Rd. #605  
Royal Oak, MI 48073  
Ph. (248)551-9100

## **Post-Operative Physical Therapy Distal Biceps Repair**

### **Post-Op:**

- Splint at 90° in supination for 2 – 3 weeks.
- Sleep with brace locked to prevent inadvertent active flexion.
- Splint / Brace
  1. Partial repair: d/c at 3 weeks
  2. Complete repair: d/c at 6 weeks
- Note angle where tension on repair – flex 40° above this, then start supination / pronation here immediately post-op.
- Begin active extension, passive flexion by 2 – 3 weeks. Limit extension to point where tension on repair noted intra-operatively.
- For motion: rest arm on table, with forearm hanging over edge. Can then actively extend. Flexion done passively.
- Sleep with brace locked to prevent inadvertent active flexion.
- Splint / Brace
  1. Partial repair: d/c at 3 weeks
  2. Complete repair: d/c at 6 weeks
- Full extension at 6 weeks.
- No active flexion until 6 weeks.
- No strengthening until 12 weeks.
- Return to vigorous labor at 6 months.
- Continue to strengthen for 1 year.